

Effects of *H pylori* infection on gap-junctional intercellular communication and proliferation of gastric epithelial cells *in vitro*

Ran Tao, Miao-Feng Hu, Jin-Tu Lou, Yong-Liang Lei

Ran Tao, Miao-Feng Hu, Jin-Tu Lou, Yong-Liang Lei, Central Laboratory, Children's Hospital, School of Medicine, Zhejiang University, Hangzhou 310003, Zhejiang Province, China
Supported by Natural Science Fund of Zhejiang Province, No. 302023

Correspondence to: Jin-Tu Lou, Central Laboratory, Children's Hospital, School of Medicine, Zhejiang University, 57 Zhugan Lane, Hangzhou 310003, Zhejiang Province, China. rtao1211@yahoo.com.cn

Telephone: +86-571-87061007-2426 Fax: +86-571-87033296

Received: May 25, 2007

Revised: August 21, 2007

communication; Gastric epithelial cell; CagA; Fluorescence redistribution after photobleaching; Methylthiazolyl tetrazolium assay

Tao R, Hu MF, Lou JT, Lei YL. Effects of *H pylori* infection on gap-junctional intercellular communication and proliferation of gastric epithelial cells *in vitro*. *World J Gastroenterol* 2007; 13(41): 5497-5500

<http://www.wjgnet.com/1007-9327/13/5497.asp>

Abstract

AIM: To explore the effects of *H pylori* infection on gap-junctional intercellular communication (GJIC) and proliferation of gastric epithelial cells *in vitro*.

METHODS: A human gastric epithelial cell line (SGC-7901) cultured on coverslips was exposed overnight to intact *H pylori* (CagA⁺ or CagA⁻ strains) and sonicated extracts, respectively. GJIC between the cells was detected by fluorescence redistribution after photobleaching (FRAP) technique. Proliferation of SGC cells was determined by methylthiazolyl tetrazolium (MTT) assay.

RESULTS: When compared with control in which cells were cultured with simple medium alone, both CagA⁺ and CagA⁻ *H pylori* isolates could inhibit GJIC (CagA⁺: $F = 57.98$, $P < 0.01$; CagA⁻: $F = 29.59$, $P < 0.01$) and proliferation (CagA⁺: $F = 42.65$, $P < 0.01$; CagA⁻: $F = 58.14$, $P < 0.01$) of SGC-7901 cells. Compared with CagA⁻ strains, CagA⁺ *H pylori* more significantly down-regulated GJIC of gastric cells (intact *H pylori*: $t = 13.86$, $P < 0.01$; sonicated extracts: $t = 11.87$, $P < 0.01$) and inhibited proliferation gastric cells to a lesser extent *in vitro* (intact *H pylori*: $t = 3.06$, $P < 0.05$; sonicated extracts: $t = 3.94$, $P < 0.01$).

CONCLUSION: Compared with CagA⁻ *H pylori* strains, CagA⁺ strains down-regulate GJIC of gastric epithelial cells more significantly and inhibit proliferation of gastric cells to a lesser extent *in vitro*. *H pylori*, especially CagA⁺ strains, may play an important role in gastric carcinogenesis.

© 2007 WJG. All rights reserved.

Key words: *H pylori*; Gap-junctional intercellular

INTRODUCTION

Epidemiological and animal studies have demonstrated a strong causal relationship between gastric cancer and chronic infection with *H pylori*, especially cytotoxin-associated gene A (*cagA*)-positive strains^[1,2]. The *cagA* gene product CagA is directly delivered into gastric epithelial cells *via* type IV secretion system. Following membrane localization and subsequent tyrosine phosphorylation, CagA interacts with a variety of host cell proteins that are involved in the regulation of cell growth and motility^[3]. However, the exact mechanism responsible for the development of gastric cancer in *H pylori*-infected patients still remains unclear.

Gap-junctional intercellular communication (GJIC) is an important mechanism controlling cellular homeostasis, proliferation and differentiation. Inhibition of GJIC between adjacent cells has been postulated to be one of the important events occurring during the promotional stage of cancer^[4]. The vast majority of neoplastic cells reduce GJIC compared to their nonneoplastic counterparts^[5]. A number of tumor promoters, such as 12-O-tetradecanoylphorbol-13-acetate (TPA), have been known as potent inhibitors of GJIC^[6].

So far, changes of GJIC in *H pylori*-associated gastric carcinoma have not been extensively exploited. In the present study, we attempted to explore the molecular mechanisms of *H pylori* infection in gastric carcinogenesis by studying its effects on GJIC of gastric epithelial cells *in vitro*.

MATERIALS AND METHODS

H pylori strains

H pylori strains 97002 and 97004 were identified by and stored in Department of Medical Microbiology and

Table 1 Effects of intact *H pylori* and sonicated extracts on GJIC of SGC-7901 cells (*n*, mean \pm SE)

Group	CagA ⁺ strain ^b	CagA ⁻ strain ^b
Intact <i>H pylori</i> ^d	26.05 \pm 3.39 (40) ^a	36.95 \pm 3.78 (44) ^a
Sonicated extracts ^d	15.92 \pm 2.53 (40) ^a	22.69 \pm 2.60 (41) ^a
Negative control	66.39 \pm 9.95 (24)	
Positive control (TPA)	8.47 \pm 0.95 (22)	

^b*P* < 0.01 one-way ANOVA vs negative control, ^a*P* < 0.05 ANOVA/Dunnett vs negative control, ^d*P* < 0.01 vs *t*-test of CagA⁺ and CagA⁻ *H pylori* strains.

Parasitology, Zhejiang University School of Medicine. The genotypes of vacuolating cytotoxin gene A (*vacA*) of the strains 97002 and 97004 were s1a/m1 and m2, respectively. The results of Western blot and cell vacuolation test demonstrated that the strain 97002 was CagA⁺/VacA⁺ and 97004 CagA⁻/VacA⁻.

H pylori culture

H pylori strains were cultured on ECY blood-free medium^[7] at 37°C for 5 d, under 100% humidity and microaerophilic conditions (50 mL/L O₂, 100 mL/L CO₂, and 850 mL/L N₂). The bacteria were harvested from the agar plates, washed twice with 0.01 mol/L PBS and stored at -20°C.

Preparation of intact *H pylori* and sonicated extract samples

The frozen bacteria were dissolved in RPMI1640 culture medium and adjusted to 1 \times 10¹⁰ CFU/L in intact bacterial samples and 1 \times 10¹² CFU/L in sonicated extract samples, respectively. The preparation of sonicated extract samples additionally included *H pylori* pulverization with ultrasound, centrifugation at 10000 r/min for 20 min with the supernatant collected.

Cell culture

Human gastric epithelial cell line SGC-7901 was obtained from Department of Medical Microbiology and Parasitology, Zhejiang University School of Medicine and cultured in RPMI1640 medium (Gibco, USA) supplemented with 10% fetal bovine serum (FBS) (Sijiqing, China), 1 \times 10⁵ IU/L penicillin and 100 mg/L streptomycin. The cells were incubated at 37°C in a humidified atmosphere containing 950 mL/L air and 50 mL/L CO₂. The cells were grown on 22 mm \times 22 mm coverslips in tissue culture dishes (35 mm in diameter) and the culture medium was changed every other day. To determine cell proliferation, SGC-7901 cells were plated into 96-well microplates (0.5 \times 10⁵ cells/well) and cultured for 12 h.

Cell treatment with *H pylori* extracts

Twenty-four hours prior to GJIC measurement, cells of the test groups were treated overnight with intact *H pylori* or sonicated extracts. Negative and positive controls were treated with RPMI1640 with 2% NBS and 5 μ g/L TPA was added to the positive control during the last 1 h.

Measurement of GJIC by FRAP technique

GJIC between SGC-7901 cells was measured by

Table 2 Effect of intact *H pylori* and sonicated extracts on proliferation of SGC-7901 cells (*n*, mean \pm SE)

Group	CagA ⁺ strain ^b	CagA ⁻ strain ^b
Intact <i>H pylori</i> ^d	0.755 \pm 0.048 (6) ^a	0.680 \pm 0.036 (6) ^a
Sonicated extracts ^d	0.938 \pm 0.037 (6)	0.830 \pm 0.056 (6) ^a
Negative control	0.955 \pm 0.038 (6)	
Positive control (TPA)	0.986 \pm 0.045 (6)	

^b*P* < 0.01 one-way ANOVA vs negative control, ^a*P* < 0.05 ANOVA/Dunnett vs negative control, ^d*P* < 0.01 vs *t*-test of CagA⁺ and CagA⁻ *H pylori* strains.

fluorescence redistribution after photobleaching (FRAP) technique first described in 1986^[8]. 6-carboxyfluorescein diacetate (6-CFDA) was used as the dye that could be retained inside the cells due to its hydrolysis by cytoplasmic esterases into 6-carboxyfluorescein (6-CF). 6-CF could permeate gap junction channels due to its low molecular weight. FRAP was achieved under a confocal laser scanning microscope (Leica TCS-SP, Germany) and the detailed protocol was performed as previously described^[9].

Determination of cell proliferation by MTT assay

When SGC-7901 cells confluent by 70% in the 96-well microplates, cell proliferation was assessed by methylthiazolyl tetrazolium (MTT) assay as previously described^[10]. The absorbance value per well at 570 nm was read on an automatic multiwell spectrophotometer (Bio-Rad, USA).

Statistical analysis

All data were presented as mean \pm SE. Statistical analysis was carried out by ANOVA followed by Dunnett's *t*-test. *P* < 0.05 was considered statistically significant.

RESULTS

H pylori down-regulated GJIC of SGC-7901 cells

The GJIC of SGC-7901 cells was measured by FRAP after treated with intact *H pylori* or sonicated extracts for 24 h and presented as fluorescence transfer rate (K, 10⁻³/s) (Table 1). In the present study, both CagA⁺ and CagA⁻ *H pylori* isolates including intact *H pylori* and sonicated extracts down-regulated GJIC of SGC-7901 cells (CagA⁺: *F* = 57.98, *P* < 0.01; CagA⁻: *F* = 29.59, *P* < 0.01). Compared with CagA⁻ strains, CagA⁺ *H pylori* more significantly down-regulated GJIC of gastric cells (intact *H pylori*: *t* = 13.86, *P* < 0.01; sonicated extracts: *t* = 11.87, *P* < 0.01). In addition, our study demonstrated that TPA (5 μ g/L for 1 h) had a significant inhibitory effect on GJIC of gastric cells.

Effect of *H pylori* on cell proliferation

The effects of intact *H pylori* and sonicated extracts on the proliferation of SGC-7901 cells were evaluated by MTT assay (A_{570 nm}) (Table 2). The results suggest that both CagA⁺ and CagA⁻ *H pylori* isolates inhibited proliferation of SGC-7901 cells (CagA⁺: *F* = 42.65, *P* < 0.01; CagA⁻: *F* = 58.14, *P* < 0.01). However, CagA⁺ *H pylori* strain inhibited proliferation of gastric cells to a lesser extent when compared with CagA⁻ strain (intact *H pylori*: *t* = 3.06, *P* < 0.05; sonicated extracts: *t* = 3.94, *P* < 0.01).

DISCUSSION

Among various forms of intercellular communication systems in multicellular organisms, GJIC is the only form by which cells exchange signals directly from the inside of one cell to the neighboring cells. GJIC plays a crucial role in maintaining homeostasis by keeping growth control signals at equilibrium among GJIC-connected cells^[11,12]. Most tumor cells have a reduced ability to communicate among themselves and/or with surrounding normal cells, confirming the importance of functional GJIC in growth control^[13-15]. GJIC is mediated by gap junction channels composed of tetramembrane spanning proteins, known as connexins. At least 13 subtypes of connexin have been identified and four or five subtypes are detectable in the gastrointestinal tract^[16].

It has been reported that connexin 32 in normal gastric mucosa is reduced significantly or absent in atrophic gastric mucosa and metaplastic epithelial cells, and no malignant cells from patients with gastric carcinoma contain detectable connexin 32^[17,18]. These results suggest that loss of cell-cell communication through the gap junction may act as an early indicator of gastric carcinoma.

In this study, the effects of *H pylori* infection on GJIC of gastric epithelial cells were detected *in vitro*, suppressing interferences of various cytokines and immune factors *in vivo*, suggesting that both CagA⁺ and CagA⁻ *H pylori* isolates inhibit GJIC of SGC-7901 cells and the down-regulating effect of CagA⁺ *H pylori* is more significant than that of CagA⁻ strains. These findings emphasize the close relationship between *H pylori* especially CagA⁺ strains and gastric carcinoma.

Increased cellular proliferation rates are characteristic in malignant tissue. Because of instability of the genome of proliferating cells, hyperproliferation increases the possibility of DNA damage and aneuploidy. Dysplasia may evolve into carcinoma if damaged DNA cannot be repaired on time or fails in promoting the apoptosis system^[19]. *H pylori* infection of the gastric mucosa is closely associated with changes in gastric epithelial cell proliferation. *In vivo* data show that gastric epithelial hyperproliferation is common in *H pylori*-infected persons and the degree of proliferation is directly associated with the severity of mucosal neutrophilic infiltration^[20-22]. However, it was reported that an overall increase in gastric epithelial cell proliferation is not associated with *H pylori* gastritis^[23]. It is not very clear whether the increased proliferation seen *in vivo* is a direct effect of *H pylori*, or a reflex increase in proliferation in response to increased cell damage, indirectly caused by *H pylori*. A recent report by Cabral *et al.*^[24] suggested that the increased cell proliferation rate in patients with *H pylori* infection might be related to the *H pylori*-induced inflammation rather than to a direct action of the pathogen.

Several *in vitro* studies reported that *H pylori* can inhibit cell proliferation^[25,26], which is consistent with the results of this study. The possible reason for the contradiction between the findings *in vivo* and *in vitro* is that *in vivo* studies are representative of the effect of persistent *H pylori* infection whereas *in vitro* experimental studies are representative of an acute *H pylori*-mediated effect. Also, the increased cell proliferation in patients with *H pylori*

infection might be due to the increased production of gastrin *in vivo*^[25]. Moreover, *in vivo* increased epithelial cell injury is associated with a reflex increase in proliferation of uninjured cells, which would not be seen *in vitro* as each cultured gastric cell is in contact with bacteria^[27]. Cell proliferation is an essential process for the integrity of gastric mucosa. Decreasing cell turnover may increase the chances of ulcer formation and delay ulcer healing. Therefore, our findings seem to be relevant to the pathogenesis of *H pylori*-associated peptic ulcer diseases.

CagA⁺ *H pylori* is frequently isolated from patients with gastric cancer in Western countries and may be more virulent in its pathogenesis^[28,29]. *In vivo* studies reported that infection with CagA⁺ *H pylori* strains is linked with higher acute inflammatory scores than CagA⁻ strains^[30,31], suggesting that these strains preferentially induce epithelial cell proliferation by stimulating inflammatory mediators. Our results show that CagA⁺ strains could inhibit proliferation of gastric epithelial cells to a lesser extent than CagA⁻ ones. Thus gastric cells injured by exposure to CagA⁺ *H pylori* strains may be more likely to progress through the cell cycle, which possibly results in the risk of replication of cells with DNA damage^[27].

In conclusion, *H pylori* can directly inhibit GJIC and proliferation of gastric epithelial cells *in vitro*. Compared with CagA⁻ *H pylori* strains, CagA⁺ strains more significantly down-regulate GJIC and inhibit proliferation to a lesser extent of gastric epithelial cells. Accelerated proliferation increases the risk of DNA damage and gene mutation. Inhibited GJIC makes cancer-initiated cells escape from the control of neighboring cells. *H pylori*, especially CagA⁺ strains, may play an important role in gastric carcinogenesis.

COMMENTS

Background

It has been widely accepted that there is a strong association between *H pylori* infection and gastric cancer, but the exact molecular mechanism of the pathogen in gastric carcinogenesis has not clarified yet. Nearly 40 years ago, loss of functional gap junctions was described in cancer cells and led to the hypothesis that such a type of intercellular communication is involved in the carcinogenesis process. Since then, a lot of data have been accumulated confirming that gap junctions are frequently decreased or absent in cancer cells. Gap junction deficiency has been defined in the literature either as the lack of gap-junction plaques or as the lack of gap-junctional intercellular communication (GJIC). It has been reported that connexin 32 in normal gastric mucosa as a mediator of GJIC is reduced significantly or absent in atrophic gastric mucosa and metaplastic epithelial cells. However, these reports have not revealed the relationship between changed GJIC and *H pylori* infection of the gastric mucosa.

Research frontiers

There has been a considerable interest over recent years in factors that predispose individuals to develop gastric carcinoma. Complex interactions between several *H pylori*, host genetics and environmental factors determine this predisposition. Understanding the molecular mechanism of the interaction between *H pylori* and gastric epithelial cells will provide us with a new strategy for effective prevention of the development of gastric cancer induced by *H pylori* infection.

Innovations and breakthroughs

In this article, the molecular mechanism of *H pylori* infection in gastric carcinogenesis was explored by studying its effects on GJIC of gastric epithelial cells *in vitro*. The results suggest that *H pylori* could inhibit GJIC of cultured gastric epithelial cells and the down-regulation effect on GJIC of CagA⁺ strains was more significant than CagA⁻ ones.

Applications

This article emphasizes the close relationship between *H pylori* especially CagA⁺ strains and gastric carcinoma. It provides a new direction to illuminate the molecular mechanism of *H pylori* in gastric carcinogenesis. It also implies that compounds able to restore GJIC in junctional deficient cells or prevent its disruption in junctional proficient cells may be used in making new strategies for the prevention and/or treatment of human gastric malignancies.

Terminology

Gap junctions: membrane structures made of intercellular channels which permit the diffusion of small hydrophilic molecules from cytoplasm to cytoplasm.

Peer review

The paper seems innovative. Altered expressions of connexins have been observed in various pathological processes of the digestive tract, including gastric cancer. To our knowledge, it is the first study to explore the molecular mechanism of *H pylori* infection in gastric carcinogenesis by studying its effects on GJIC of gastric epithelial cells *in vitro*.

REFERENCES

- Kelley JR, Duggan JM. Gastric cancer epidemiology and risk factors. *J Clin Epidemiol* 2003; **56**: 1-9
- Zheng Q, Chen XY, Shi Y, Xiao SD. Development of gastric adenocarcinoma in Mongolian gerbils after long-term infection with *Helicobacter pylori*. *J Gastroenterol Hepatol* 2004; **19**: 1192-1198
- Hatakeyama M. The role of *Helicobacter pylori* CagA in gastric carcinogenesis. *Int J Hematol* 2006; **84**: 301-308
- Kang KS, Yun JW, Yoon B, Lim YK, Lee YS. Preventive effect of germanium dioxide on the inhibition of gap junctional intercellular communication by TPA. *Cancer Lett* 2001; **166**: 147-153
- Trosko JE, Ruch RJ. Cell-cell communication in carcinogenesis. *Front Biosci* 1998; **3**: d208-d236
- Ruch RJ, Trosko JE, Madhukar BV. Inhibition of connexin43 gap junctional intercellular communication by TPA requires ERK activation. *J Cell Biochem* 2001; **83**: 163-169
- Fang PC, Zhu YL, Yin X, Wu QD, Lan MG, Wu PJ. Study on ECY blood-free medium for the isolation of *Helicobacter pylori*. *Zhonghua Yixue Jianshan Zazhi* 1993; **16**: 131-133
- Wade MH, Trosko JE, Schindler M. A fluorescence photobleaching assay of gap junction-mediated communication between human cells. *Science* 1986; **232**: 525-528
- Mao GG, Fu YT, Ye SJ. Determination of gap junctional intercellular communication in cultured cells. *Zhonghua Laodong Weisheng Zhiyebing Zazhi* 2000; **18**: 376-377
- Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983; **65**: 55-63
- Yamasaki H. Role of disrupted gap junctional intercellular communication in detection and characterization of carcinogens. *Mutat Res* 1996; **365**: 91-105
- Yamasaki H, Krutovskikh V, Mesnil M, Tanaka T, Zaidan-Dagli ML, Omori Y. Role of connexin (gap junction) genes in cell growth control and carcinogenesis. *C R Acad Sci III* 1999; **322**: 151-159
- Yamasaki H, Omori Y, Zaidan-Dagli ML, Mironov N, Mesnil M, Krutovskikh V. Genetic and epigenetic changes of intercellular communication genes during multistage carcinogenesis. *Cancer Detect Prev* 1999; **23**: 273-279
- Ruch RJ, Porter S, Koffler LD, Dwyer-Nield LD, Malkinson AM. Defective gap junctional intercellular communication in lung cancer: loss of an important mediator of tissue homeostasis and phenotypic regulation. *Exp Lung Res* 2001; **27**: 231-243
- Trosko JE, Chang CC. Role of stem cells and gap junctional intercellular communication in human carcinogenesis. *Radiat Res* 2001; **155**: 175-180
- Nishitani A, Hirota S, Nishida T, Isozaki K, Hashimoto K, Nakagomi N, Matsuda H. Differential expression of connexin 43 in gastrointestinal stromal tumours of gastric and small intestinal origin. *J Pathol* 2005; **206**: 377-382
- Uchida Y, Matsuda K, Sasahara K, Kawabata H, Nishioka M. Immunohistochemistry of gap junctions in normal and diseased gastric mucosa of humans. *Gastroenterology* 1995; **109**: 1492-1496
- Nagahara A, Watanabe S, Miwa H, Endo K, Hirose M, Sato N. Reduction of gap junction protein connexin 32 in rat atrophic gastric mucosa as an early event in carcinogenesis. *J Gastroenterol* 1996; **31**: 491-497
- Gao H, Wang JY, Shen XZ, Liu JJ. Effect of *Helicobacter pylori* infection on gastric epithelial cell proliferation. *World J Gastroenterol* 2000; **6**: 442-444
- Fraser AG, Sim R, Sankey EA, Dhillon AP, Pounder RE. Effect of eradication of *Helicobacter pylori* on gastric epithelial cell proliferation. *Aliment Pharmacol Ther* 1994; **8**: 167-173
- Bechi P, Balzi M, Becciolini A, Maugeri A, Raggi CC, Amorosi A, Dei R. *Helicobacter pylori* and cell proliferation of the gastric mucosa: possible implications for gastric carcinogenesis. *Am J Gastroenterol* 1996; **91**: 271-276
- Murakami K, Fujioka T, Kodama R, Kubota T, Tokieda M, Nasu M. *Helicobacter pylori* infection accelerates human gastric mucosal cell proliferation. *J Gastroenterol* 1997; **32**: 184-188
- Chow KW, Bank S, Ahn J, Roberts J, Blumstein M, Kranz V. *Helicobacter pylori* infection does not increase gastric antrum mucosal cell proliferation. *Am J Gastroenterol* 1995; **90**: 64-66
- Cabral MM, Oliveira CA, Mendes CM, Guerra J, Queiroz DM, Rocha GA, Rocha AM, Nogueira AM. Gastric epithelial cell proliferation and cagA status in *Helicobacter pylori* gastritis at different gastric sites. *Scand J Gastroenterol* 2007; **42**: 545-554
- Ricci V, Ciacci C, Zarrilli R, Sommi P, Tummuru MK, Del Vecchio Blanco C, Bruni CB, Cover TL, Blaser MJ, Romano M. Effect of *Helicobacter pylori* on gastric epithelial cell migration and proliferation in vitro: role of VacA and CagA. *Infect Immun* 1996; **64**: 2829-2833
- Pearce HR, Kalia N, Bardhan KD, Atherton JC, Brown NJ. Effects of *Helicobacter pylori* on endothelial cell proliferation and chemotaxis. *Digestion* 2004; **69**: 201-210
- Smoot DT, Wynn Z, Elliott TB, Allen CR, Mekasha G, Naab T, Ashktorab H. Effects of *Helicobacter pylori* on proliferation of gastric epithelial cells in vitro. *Am J Gastroenterol* 1999; **94**: 1508-1511
- Blaser MJ, Perez-Perez GI, Kleanthous H, Cover TL, Peek RM, Chyou PH, Stemmermann GN, Nomura A. Infection with *Helicobacter pylori* strains possessing cagA is associated with an increased risk of developing adenocarcinoma of the stomach. *Cancer Res* 1995; **55**: 2111-2115
- Hatakeyama M, Higashi H. *Helicobacter pylori* CagA: a new paradigm for bacterial carcinogenesis. *Cancer Sci* 2005; **96**: 835-843
- Peek RM Jr, Miller GG, Tham KT, Perez-Perez GI, Zhao X, Atherton JC, Blaser MJ. Heightened inflammatory response and cytokine expression in vivo to cagA⁺ *Helicobacter pylori* strains. *Lab Invest* 1995; **73**: 760-770
- Bhat N, Gaensbauer J, Peek RM, Bloch K, Tham KT, Blaser MJ, Perez-Perez G. Local and systemic immune and inflammatory responses to *Helicobacter pylori* strains. *Clin Diagn Lab Immunol* 2005; **12**: 1393-1400

S- Editor Liu Y L- Editor Wang XL E- Editor Yin DH